

DISCRETE MATHEMATICS PROBLEMS. JUNE 20, 2002

INSTRUCTOR: LÁSZLÓ BABAI

Exercise 1. Given $B \in \mathbb{R}^{n \times n}$ and $x \in \mathbb{R}^n$, show that if $(\forall x)(xBx^t = 0)$ and $B = B^t$, then $B = 0$.

Exercise 2. If A is an $n \times n$ orthogonal matrix (i.e., $AA^t = I$) and λ is an eigenvalue of A , then $|\lambda| = 1$.

Exercise 3. Prove from first principles (i.e., without using the determinant) that the row rank of a matrix equals the column rank. *Hint:* Gaussian elimination.

Definition 4. The determinant rank of a matrix is the size of the largest non-singular submatrix. A square matrix is *non-singular* if its determinant is non-zero.

Exercise 5. Prove that the determinant rank of a matrix equals the row rank, which equals the column rank.

Recall Lindsey's lemma: If A is an $a \times b$ submatrix of a Hadamard matrix, then $|\sum a_{ij}| \leq \sqrt{abn}$.

Exercise 6. It was shown in class that the payoff for the Gale-Berlekamp switching game is $\leq \sqrt{2}n^{3/2}$. Replace this with $\leq (1 + o(1))n^{3/2}$.

Hint: Prime Number Theorem. Note: $a_n = o(1)$ is equivalent to $a_n \rightarrow 0$.

Exercise 7. We want to pick numbers from $1, \dots, n$ such that there is no 3-term arithmetic progression among the numbers selected. In class we gave an algorithm for doing this and getting $\Omega(\sqrt{n})$ numbers; now find an *explicit* subset of this size.

Definition 8. Given a k -colored matrix (i.e., one with entries $1, \dots, k$), a homogeneous submatrix is a submatrix all of whose entries are equal.

Exercise 9. Show that $(\forall k, \ell)(\exists n_0)$ such that if $n > n_0$, then every k -colored $n \times n$ matrix has a homogeneous $\ell \times \ell$ submatrix.

Prove this for $\ell = \Omega\left(\frac{\log n}{\log k}\right)$.

Hint: Focus on $k = 2$.

Exercise 10. As above, let $k = 2$, and show that $\ell = O(\log n)$; i.e., there exist $n \times n$ $(0, 1)$ -matrices such that every homogeneous $\ell \times \ell$ submatrix satisfies $\ell = O(\log n)$.

Hint: Probabilistic method. (Flip coins to decide the entries.)

Exercise 11. Prove: if an $n \times n$ Hadamard matrix has an $\ell \times \ell$ homogeneous submatrix, then $\ell \leq \sqrt{n}$.

Hint: Lindsey's Lemma

OPEN QUESTION. Construct an explicit family of $n \times n$ (± 1) matrices such that all $\ell \times \ell$ homogeneous submatrices have $\ell \leq n^{49}$. A candidate is the Quadratic Residue matrix over \mathbb{F}_p , defined by $a_{i,j} = \left(\frac{i+j}{p}\right)$.

Exercise 12. If q is a square ($q = p^{2k}$), then the Quadratic Residue matrix over \mathbb{F}_q *does* have a homogeneous submatrix of size $\sqrt{q} \times \sqrt{q}$. Since Weil's Theorem does not distinguish between square and non-square prime powers, this suggests that it cannot be used to resolve the Open Question above.

Exercise 13. The Sylvester matrix S_k (of size $2^k \times 2^k$) has $\sqrt{n} \times \sqrt{n}$ homogeneous submatrices.