

Discrete Math, 22th day, Friday 8/13/04  
REU 2004. Info:  
<http://people.cs.uchicago.edu/~laci/reu04>.

Instructor: László Babai  
Scribe: Charilaos Skiadas

## 1 0, 1-measures

**Exercise 22.1.** Prove the following theorem. (*Hint:* Use Zorn's Lemma.)

**Theorem 22.2.** *If  $\Omega$  is an infinite set, then there exists a finite additive nontrivial 0, 1 measure on  $2^\Omega$ ,  $\mu : 2^\Omega \rightarrow \{0, 1\}$ , such that  $\mu(\Omega) = 1$ . Nontrivial means that  $\mu(\{a\}) = 0$  for all  $a$ .*

This is a hint to the infinite switch problem.

## 2 Sign-Rigidity (Paturi-Simon)

Let  $M$  be a *sign matrix*, i.e., a matrix consisting of  $\pm 1$  entries. A matrix  $A$  of the same dimensions as  $M$  *realizes*  $M$  if the sign of  $a_{i,j}$  is equal to  $m_{i,j}$ . The *sign-rank* of  $M$  is the minimum rank of a matrix realizing  $M$ .

**Exercise 22.3 (Alon-Frankl-Rödl, 1984).** Use Warren's theorem to prove that almost all  $n \times n$  sign-matrices have sign-rank greater than or equal to  $\frac{n}{32}$ .

Finding explicit matrices that satisfy the preceding example is a hard problem. No particular examples are known.

**Conjecture 22.4.** *Hadamard matrices have sign-rank  $\geq cn$ .*

For applications, all we need would be an explicit matrix with rank greater than  $n^\epsilon$  for some fixed  $\epsilon > 0$ . The best known explicit bound was  $\Omega(\log n)$  until 2002.

**Theorem 22.5 (Forster 2002).** *Let  $X \subseteq \mathbb{R}^k$  such that  $|X| \geq k$  and the elements of  $X$  are in general position (i.e.,  $k$ -wise linearly independent). Then there exists an invertible  $k \times k$  matrix  $A$  such that  $\sum_{x \in X} \frac{(Ax)(Ax)^T}{(Ax)^T(Ax)} = aI_k$ , a constant multiple of the identity matrix.*

If we assume that the theorem is true, the value of  $a$  can easily be found by taking the trace on both sides (recall that the trace of the product of two matrices is independent of the order). This gives  $ak = \sum_{x \in X} 1 = |X|$

**Exercise 22.6 (Forster).** If  $A$  realizes  $M$ , then  $\text{rk}(A) \geq \frac{n}{\|M\|}$ . Recall that  $\|M\| = \max_{\|x\|=1} \|Mx\| = \sqrt{\lambda_{\max}(M^T M)}$  by the spectral theorem.

**Corollary 22.7.** If  $H$  is a Hadamard matrix, then  $\text{sign-rank}(H) \geq \sqrt{n}$ .

**Proof:** If  $A$  realizes  $H$ , then  $\text{rk}(A) \geq \frac{n}{\|H\|}$ . Since  $H^T H = nI$ , we get that  $\|H\|^2 = \lambda_{\max}(nI) = n$